

Study on responses of C-peptide and blood glucose for Carbohydrate 70g in diabetes

Hiroshi Bando^{1,2*}, Koji Ebe^{2,3}, Tetsuo Muneta^{2,4}, Masahiro Bando⁵ and Yoshikazu Yonei⁶

¹Tokushima University/Medical Research, Tokushima, Japan

²Japan Low Carbohydrate Diet Promotion Association, Kyoto, Japan

³Takao Hospital, Kyoto, Japan

⁴Muneta Maternity Clinic, Chiba, Japan

⁵Department of Nutrition and Metabolism, Institute of Biomedical Sciences, Tokushima University Graduate School, Tokushima, Japan

⁶Anti-Aging Medical Research Center, Graduate School of Life and Medical Sciences, Doshisha University, Kyoto, Japan

*Corresponding author: Hiroshi Bando, MD, Tokushima University /Medical Research Nakashowa 1-61, Tokushima 770-0943 Japan, Tel: +81-90-3187-2485; E-mail: pianomed@bronze.ocn.ne.jp

Received Date: 25 June 2018; Accepted Date: 27 July 2018; Published Date: 08 August 2018

Copyright: © 2018 Bando H, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Bando H, Ebe K, Muneta T, Bando M, Yonei Y (2018) Study on responses of C-peptide and blood glucose for Carbohydrate 70g in diabetes. Endocrinol Metab Vol. 2 No.2: 110

Abstract

Background: Discussion concerning Carbohydrate Restriction (CR) and Low Carbohydrate Diet (LCD) has been continued. Authors formerly proposed insulinogenic index-carbohydrate 70 g (IGI-carbo70). Along this, C-peptide index for Carbohydrate-70 g (CPI-Carbo70) is studied.

Subjects and Methods: Subjects were 46 patients (62.5 yo. in average) with Type 2 diabetes mellitus (T2DM) and admitted for evaluation. CR diet was given on day 1-2 with 60% carbohydrates, 25% lipids and 15% protein with 1400 kcal/day. On day 2, breakfast with Carbo70 was given, and glucose and C-peptide (CP) were measured at 0 and 30 minutes and analyzed.

Results: As basal data, average HbA1c was 8.1%, median HbA1c, average daily glucose, median M value was 7.7%, 187 mg/dL, 71, respectively. In response to Carbo70, median values of glucose-0, glucose-30, delta-glucose, CP-0, CP-30, delta-CP was 142, 186, 22 mg/dL, 0.9, 1.4, 0.4 ng/mL, respectively. CPI-Carbo70 was 1.54 in median. There were significant correlation between glucose and CP on 0, 30min, delta-0-30min ($p < 0.01$). Significant correlation was observed between CPI and HbA1c, CPI and M value ($p < 0.01$).

Discussion and Conclusion: CPI-Carbo70 might be useful for evaluating pancreas function and glucose variability in clinical research. Current data would become the fundamental data for the future diabetic practice and research development.

Keywords: C-peptide (CP); C-peptide index for Carbohydrate-70 (CPI-Carbo70); Insulinogenic index (IGI); low carbohydrate diet (LCD); type 2 diabetes mellitus (T2DM); Calorie Restriction (CR)

Abbreviations: CR: Carbohydrate Restriction; LCD: Low Carbohydrate Diet; CPI-Carbo70: C-peptide index for Carbohydrate-70 g; T2DM: Type 2 diabetes mellitus; ADA: American Diabetes Association; ACP: American College of Physicians; IDF: International Diabetes Federation; 75 g OGTT: 75 g oral glucose tolerance test; IGI-Carbo70: insulinogenic index (IGI)-Carbohydrate70 g; M: Morbus value; MAGE: mean amplitude of glycemic excursions.

Introduction

Metabolic syndrome (Met-S) has been prevalent in the world nowadays. It was proposed and investigated from syndrome X in 1980s associated with insulin resistance [1]. Met-S has been gradually increased with medical, social and economic problems [2], and it is also defined as a cluster of mutual metabolic abnormalities accompanied by glucose metabolism, including dyslipidemia, elevated blood pressure and central obesity [3].

As mentioned above, impaired glucose metabolism would exist in the fundamental pathophysiology of Met-S. Patients with diabetes mellitus have been increasing worldwide [4]. There was recently the changes in the guideline concerning diabetes. American Diabetes Association (ADA) gave comments in 2017 [5], which was followed by the joint algorithm of European Diabetes Society (EASD) 2012 [6]. After that, American College of Physicians (ACP) proposed the change in standard value about HbA1c [7], in which the

management goal for HbA1c in most type 2 diabetic patients would be 7% or more and less than 8%. Against the concept of ACP, ADA made an objection comment immediately [8]. Thus, diabetic management has been in discussion among several guidelines from medical societies, leading to better clinical practice with accumulated evidences.

Moreover, there was a large-scale epidemiological study, which is "The Prospective Urban Rural Epidemiology (PURE) study". It included wide data from 140 thousands subjects from 17-18 countries in the world [9]. The result revealed that there were higher carbohydrate intake associated with an increased risk of HR 1.28 of total mortality. Then, it seems to be recommended to take lower carbohydrate for everyone to lead a better life [10].

International Diabetes Federation (IDF) summarized and proposed the Standards of Medical Care in Diabetes [11]. Among this guideline, the crucial points seemed to be monitoring carbohydrate intake, carbohydrate counting and experience-based estimation. In the future, continuing evaluation and investigation of eating patterns and macronutrient distribution would be beneficial for achieving better glycemic control.

Regarding the nutritional treatment for diabetes mellitus, there has been continuing discussion the comparison about Calorie Restriction (CR) diet and Low Carbohydrate Diet (LCD) [12,13]. Originally, LCD was initiated by Bernstein who was physician with type 1 diabetes mellitus (T1DM). After that, LCD became rather popular and known widely [14]. In Japan, authors and colleagues have started LCD and developed medically and socially with Japan LCD promotion association, and have given adequate therapy for many diabetic patients with successful treatment [15]. Furthermore, we have continued LCD research related to lipid metabolism, renal function, elevated ketone bodies and three types of LCD formula which are petit, standard and super LCD [16-18].

As we have continued clinical research about CR /LCD, and related investigation, we have proposed a trial of clinically new index which is simple and useful method. It is similar procedure and calculation of insulinogenic index (IGI) to 75 g oral glucose tolerance test (75 g OGTT). Subjects have breakfast including 70 g of carbohydrate, and blood samples are drawn before and 30 minutes after for the measurement of blood glucose and immunoreactive insulin (IRI) [19]. It is called insulinogenic index (IGI)-Carbohydrate 70 g (IGI-Carbo70), and seems to become useful for clinical diabetic practice.

We develop this evaluation method, try to use the measurement of C-Peptide (CP) value instead of IRI and report the data in this study.

Subjects and Methods

In current study, the subjects were 46 patients (M/F 19/27) enrolled with Type 2 diabetes mellitus (T2DM). Fundamental data were shown in Table 1. Average age is 62.5 +/- 10.9 years old (mean +/- standard deviation) and 65 (55- 68) years old

(median [25-75%]). They were admitted to the hospital for 14 days, which purpose would be further evaluation, examination and treatment. They were in-patient and on our protocol of CR and LCD nutritional therapy.

Methods are described in the following: 1) Formula diet for CR was given to the subjects on day 1 and 2. CR includes 60% carbohydrates, 25% lipids and 15% protein with 1400 kcal/day. 2) Formula meal of super LCD was provided from 3 to 14 days, including 12% carbohydrates, 64% lipids and 24% protein with 1400 kcal/day.

This protocol with CR and LCD has been continued in our clinical research for years. However, this study has only indicated for the breakfast on day 2. Formula CR meal has 840 kcal of carbohydrate per day including 210g of carbohydrate totally in 3 meals. According to the guideline of Japan Diabetes Society (JDS), standard nutrients balance of PFC (protein, fat, carbohydrate) are 60% carbohydrates, 25% lipids and 15% protein [20]. In this manner, formula breakfast including 70g of carbohydrate was provided after overnight fasting.

Detail protocol is shown as follows: 1) Basal biomarkers were checked in fasting situation on day 2, 2) blood glucose level and C-peptide value were measured before breakfast (0 min), 3) patient eats breakfast with 70 g of carbohydrate, 4) levels of glucose and C-peptide were measured 30 minutes after breakfast, 5) both markers on 0, 30 min and increment of glucose and C-peptide were measured, 6) the index of ratio for increment of C-peptide / glucose was calculated as CPI-Carbo70, 7) The detail calculation method is as follows: Increment is described by delta, delta-glucose and delta-CP are calculated, delta-CP/delta-glucose x 100 equals CPI-Carbo70.

Morbus value and glucose profile

As to the daily glucose profile on day 2, blood glucose was measured 7 times a day, which time was 8, 10, 12, 14, 17, 19, 22 h. From these data, average glucose level and Morbus (M) value were calculated.

M value has been known the useful index which indicates both blood sugar level and mean amplitude of glycemic excursions (MAGE) [21,22]. From the glucose profiles, average glucose level and M value could be calculated. M value is useful for supposing the status of MAGE. M value is characteristic for a logarithmic transformation of the deviation of glycemia from an arbitrary assigned "ideal" glucose value. It includes the degree of mean glucose value and glucose swings [21-23].

The definition of M value has been as follows: $M = MBS + MW$, where $MW = (\text{maximum blood glucose} - \text{minimum glucose}) / 20$; $MBS = \text{the mean of MBSBS}$; $MBSBS = \text{individual M-value for each blood glucose value calculated as } (\text{absolute value of } [10 \times \log (\text{blood glucose value} / 120)])^3$ [21-23].

The significance and interpretation of M value has been in the following: the standard range is <180, borderline is 180-320 and abnormal range is >320. According to some experimental research, similar results were taken for measuring 7 times or 20 times a day [22- 24]. Furthermore, its

revealed similar result compared with continuous glucose monitoring (CGM) [22,24,25].

Statistical analyses

In this study, data were indicated as the mean \pm standard deviation (SD) and also as median, quartile of 25% and 75% according to the biomarkers. In statistical analyses, correlation coefficients were calculated by using Spearman test of the Microsoft Excel analytical tool, which is Four steps Excel Statistics 4th edition [26].

Ethical considerations

This investigation was conducted in compliance with the ethical principles of the Declaration of Helsinki. It was also conducted with Japan's Act on the Protection of Personal Information along with the Ministerial Ordinance on Good Clinical Practice (GCP) for Drug (Ordinance of Ministry of Health and Welfare No. 28 of March 27, 1997).

In addition, an ethical committee was established including doctor, nurse, pharmacist and expert in the medical and legal specialty. We have discussed enough and got the conclusion that this study has been valid and agreed without any problems. Furthermore, informed consents and written paper agreements have been obtained from the all subjects. The study has been registered with UMIN #R000031211.

Results

Basal data

Current study enrolled 46 patients with T2DM. Fundamental diabetic data were shown in **Table 1**. Average glucose level was 205 mg/dL and median M value was 71.

Table 1 Subjects and basal data.

	Mean \pm SD	Median [25-75%]
Subjects number (M/F)	46 (19/27)	46 (19/27)
age (years old)	62.5 \pm 10.9	65 [55-68]
Glucose profile		
HbA1c (%)	8.1 \pm 1.8	7.7 [7.0-9.2]
average glucose (mg/dL)	204 \pm 71.2	187 [161-228]
Morbus value	187 \pm 256	71 [43-203]

C-peptide index for Carbo70

Responses of blood glucose and C-peptide (CP) for carbohydrate 70 g were shown in **Table 2**. The values are expressed by average \pm standard deviation and median [25%-75%]. As median value, delta-glucose, delta-CP, CPI-Carbo70 was 22 mg/dL, 0.4 ng/mL, 1.54, respectively.

Table 2 Responses of Glucose and C-peptide for Carbo-70.

	Mean \pm SD	Median [25-75%]
Response of blood glucose before (0 min)	163 \pm 54.9	142 [134-186]
after (30 min)	195 \pm 65.2	186 [155-214]
increment (Δ)	322 \pm 26.4	22 [13-46]
Response of C-peptide before (0 min)	1.08 \pm 0.65	0.9 [0.7-1.4]
after (30 min)	1.52 \pm 0.83	1.4 [1.0-1.9]
increment (Δ)	0.44 \pm 0.41	0.4 [0.8-2.5]
Index of CRP for carbo-70		
ACPR/ABG	1.94 \pm 1.51	1.54 [0.8-2.5]

Correlation among biomarkers

There was significant correlation between fasting glucose and C-peptide ($p < 0.01$) (**Figure 1**). In response to the loading of carbohydrate 70g, blood glucose and C-peptide were measured at 30 min. There was significant correlation between glucose and C-peptide on 30 min ($p < 0.01$) (**Figure 2**).




Figure 1 Correlation between Fasting glucose and C-peptide.




Figure 2 Correlation between glucose and C-peptide on 30 min.

C-peptide Index (CPI) is calculated by delta-CP/delta-glucose \times 100. There was significant correlation between delta-CP and delta-glucose ($p < 0.01$) (**Figure 3**).




Figure 3 Correlation between delta-glucose and delta-C-peptide on 0-30 min.

Regarding CPI, two correlations were investigated. Significant negative correlation was observed between CPI and HbA1c ($p < 0.01$) (**Figure 4**). Furthermore, significant negative correlation was observed between CPI and M value ($p < 0.01$) (**Figure 5**).




Figure 4 Correlation between HbA1c and CPI-Carbo70.




Figure 5 Correlation between M value and CPI-Carbo70.

Discussion

Authors and colleagues have continued clinical practice and research concerning LCD for years [17,18,27]. Furthermore, we have developed social movement for healthier life through Japan Low Carbohydrate Diet Promotion Association.

In this study, 70 g of Carbohydrate as a breakfast was used. **Figures 1-3** revealed 3 significant correlations between glucose and C-peptide in 0 min, 30 min, and delta-0-30min. These high correlation seems to suggest the validity and usefulness of this protocol. These statistical precise data in this order (0 min, 30

min, delta-0-30min) was $r = 0.395$ / $p = 0.0078$, $r = 0.431$ / $p = 0.0038$, $r = 0.510$ / $p = 0.0005$, respectively. Among these, delta-0-30min showed higher correlation degree, supposing that delta-glucose and delta-CP may be useful and meaningful in clinical practice and research.

Furthermore, **Figures 4 and 5** revealed significant correlations between CPI-Carbo70 and HbA1c, CPI-Carbo70 and M value, in which the latter revealed higher degree of correlation. When compared the difference of numerical values in HbA1c and M value, the latter has larger difference result than that of the former. Consequently, it may be more useful by M value for analyzing biomarkers in related research.

In relation to current study, there have been several reports concerning the responses of glucose, insulin and C-peptide against carbohydrate loading on breakfast.

There was a study of postprandial hyperglycemia (PPHG) after lunch and dinner in the case of with and without breakfast [28]. The result was that skipping breakfast increases PPHG with impaired insulin response and lower incremental GLP-1. It may suggest the influence of breakfast on glucose regulation that persists throughout the day. Without breakfast, AUC of blood glucose (0-30 min) of lunch increased by 23.7%, while AUC of C-peptide decreased by 42.5% after lunch. However, basal value of C-peptide was 3.7 ng/mL and 1.8 ng/mL, with breakfast and without breakfast, respectively. Consequently, it seems to be rather difficult to compare the both and to speculate the detail.

In other study, T2DM patients were investigated with 2 kinds of breakfast provided [29]. One is carbo-breakfast with PFC=15:20:65%, another is protein-breakfast with PFC=35:20:45%. When lunch including 65% of carbohydrate was given to both groups, protein-group showed higher insulin and C-peptide response, with lower postprandial hyperglycemia. Insulin, C-peptide and GIP levels showed the second-meal phenomenon, keeping the glucose concentrations controlled.

There was also an experiment for glucose increase in four conditions on ingestion of carbohydrate 50 g [30]. These protocols are as follows: only rice, salad + rice, rice + yogurt, and yogurt + rice, respectively. The result for the increment of blood glucose on 30 minute was 39.0 mg/dL, 19.4 mg/dL, 29.2 mg/dL, 15.9 mg/dL, respectively. The latter three group revealed reduced postprandial hyperglycemia compared with only rice group. These results would suggest the effect of taking salad and/or yogurt together for decreasing the risk of high glucose.

Furthermore, some reports were found with mixed meal challenge on breakfast. T2DM patients were provided 3 kinds of breakfast, which are 1) WB: whey diet 28 g with 42 g total protein, 2) PB diet: various protein 42 g, 3) CB: high-carbo with 17 g protein [31]. Overall postprandial AUC glucose was reduced by 12% in PB diet and by 19% in WB diet, compared with CB diet. Increment of C-peptide would be 7.4, 4.0, 1.0 pg/mL, respectively. After 12 weeks, decrease of HbA1c and body weight were, 0.89%, 0.6%, 0.36%, and 7.6kg, 6.1 kg, 3.5

kg, respectively. Thus, whey protein-based breakfast could become an important adjuvant for T2DM.

Responses of glucose, insulin, C-peptide and Triglyceride were investigated between Red Meat/Refined Grain Diet and Dairy/Chicken/Nuts/Whole Grain Diet [32]. In each group, carbohydrate amount per day and CP-incremental AUC were 218 g vs 181 g, 8.6 vs 6.4 nmol/L/3 h ($p < 0.002$), respectively. Compared the results from insulin and c-peptide data, insulin sensitivity and hepatic insulin clearance would be involved.

In response to a mixed meal challenge, GLP-1-induced insulin secretion was found [33]. It was significantly reduced in people with impaired glucose tolerance, compared with those with normal glucose tolerance.

After mixed nutrient preload, glucose tolerance was relieved in proportion to the degree of T2DM [34]. Investigating responses to nutrient ingestion might reveal pathophysiological mechanism of T2DM, leading to a potent tool to improve glycemic control.

As mentioned above, researches on breakfast containing complex nutrients are progressing. Since the content is not single nutrient, these studies would have limits and problems to be solved. They may include various kinds of mixture for nutrients in the meal, different speed of digestion and absorption for each subjects, and response ability of insulin and C-peptide secretion. However, our current research concerning CPI-Carbo70 might become a reference data for future diabetic practice and research.

Conclusion

In current study, we investigated the response of blood glucose and C-peptide for carbohydrate 70 g and proposed the usefulness of CPI-Carbo70. This index could become one of the simple and useful tool for evaluating pancreas function and glucose variability. This report would provide one of the basal data in the clinical practice and research development.

Acknowledgement

The part of the content of this article was presented at the 90th Scientific Meeting of Japan Diabetes Society (JDS) Annual Congress, Tokyo, 2018. The authors would like to thank all patients and staffs for their understanding and cooperation.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

References

1. Reaven GM (1988) Role of insulin resistance in human disease. *Diabetes* 37: 1595-607.
2. McGill AT (2014) Past and future corollaries of theories on causes of metabolic syndrome and obesity related comorbidities part 2: a composite unifying theory review of human-specific co-adaptations to brain energy consumption. *Arch Public Health* 72: 31.
3. Amihaesei IC, Chelaru L (2014) Metabolic syndrome a widespread threatening condition; risk factors, diagnostic criteria, therapeutic options, prevention and controversies: an overview. *Rev Med Chir Soc Med Nat Iasi* 118: 896-900.
4. World Health Organization 2016. Global report on diabetes. ISBN 978 92 4 156525 7 (NLM classification WK 810).
5. American Diabetes Association (2017) Pharmacologic Approaches to Glycemic Treatment. *Diabetes Care* 40: S64-S74.
6. Giaccari A, Giorda CB, Riccardi G, De Micheli A, Bruno G, et al. (2012) Management of hyperglycemia in type 2 diabetes: a patient-centered approach. Position statement of the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care* 35: 1364-1379.
7. American College of Physicians (2017) Clinical Guidelines and Recommendations. Web site.
8. American Diabetes Association (2018) Pharmacologic Approaches to Glycemic Treatment: Standards of Medical Care in Diabetes-2018. *Diabetes Care* 41: S73-S85.
9. Miller V, Yusuf S, Chow CK, Dehghan M, Corsi DJ, et al. (2016) Availability, affordability, and consumption of fruits and vegetables in 18 countries across income levels: findings from the Prospective Urban Rural Epidemiology (PURE) study. *Lancet Glob Health* 4: e695-703.
10. Dehghan M, Mente A, Zhang X, Swaminathan S, Li W, et al. (2017) Prospective Urban Rural Epidemiology (PURE) study investigators (2017) Associations of fats and carbohydrate intake with cardiovascular disease and mortality in 18 countries from five continents (PURE): a prospective cohort study. *Lancet* 390: 2050-2062.
11. Ogurtsova K, da Rocha Fernandes JD, Huang Y, Linnenkamp U, Guariguata L, et al. (2017) IDF Diabetes Atlas: Global estimates for the prevalence of diabetes for 2015 and 2040. *Diabetes Res Clin Pract* 128: 40-50.
12. Shai I, Schwarzfuchs D, Henkin Y, Shahar DR, Witkow S, et al. (2008) Weight loss with a low-carbohydrate, mediterranean, or low-fat diet. *N Engl J Med* 359: 2292-2303.
13. Atallah R, Fillion KB, Wakil SM, Genest J, Joseph L, et al. (2014) Long-Term Effects of 4 Popular Diets on Weight Loss and Cardiovascular Risk Factors: A Systematic Review of Randomized Controlled Trials. *Circ Cardiovasc Qual Outcomes* 7: 815-827.
14. Bernstein RK (2007) Dr. Bernstein's Diabetes solution: The Complete Guide to Achieving Normal Blood Sugars. Little, Brown US, New York.
15. Bando H, Ebe K, Nakamura T, Bando M, Yonei Y (2016) Low Carbohydrate Diet (LCD): Long and short-term effects and hyperketonemia. *Glycative Stress Res* 3: 193-204.
16. Muneta T, Kawaguchi E, Nagai Y, Matsumoto M, Ebe K, et al. (2016) Ketone body elevation in placenta, umbilical cord, newborn and mother in normal delivery. *Glycative Stress Res* 3: 133-140.
17. Bando H, Ebe K, Muneta T, Bando M, Yonei Y (2017) Investigation of uric acid and cystatin C on low-carbohydrate diet (LCD). *Diabetes Res Open J* 2017; 3: 31-38.
18. Ebe K, Bando H, Muneta T, Bando M, Yonei Y (2017) Effect of low carbohydrate diet (LCD) for diabetic patients with hypertriglyceridemia. *Endocrinol Metab* 1: 104.

19. Bando H, Ebe K, Muneta T, Bando M, Yonei Y (2017) Proposal for Insulinogenic Index (IGI)-Carbo70 as Experimental Evaluation for Diabetes. *J Clin Exp Endocrinol* 1: 102.
20. Japan Diabetes Association (2013) Diabetes clinical practice guidelines Based on scientific evidence.
21. Schlichtkrull J, Munck O, Jersild M (1965) The M-value, an index of blood sugar control in diabetics. *Acta Med Scand* 177: 95-102.
22. Siegelaar SE, Holleman F, Hoekstra JBL, Devries JH (2010) Glucose Variability; Does It Matter? *Endocrine Reviews* 31: 171-182.
23. Service FH (2013) Glucose variability. *Diabetes* 62: 1398-1404.
24. Monnier L, Colette C (2011) Glycemic Variability: Can We Bridge the Divide Between Controversies? *Diabetes Care* 34: 1058-1059.
25. Baghurst P (2011) Calculating the mean amplitude of glycemic excursion from continuous glucose monitoring data: an automated algorithm. *Diabetes Technol Ther* 13: 296-302.
26. Yanai H (2015) Four step excel statistics, 4th Edition, Seiun-sha Publishing Co.Ltd, Tokyo.
27. Ebe K, Bando H, Yamamoto K, Bando M, Yonei Y (2018) Daily carbohydrate intake correlates with HbA1c in low carbohydrate diet (LCD). *J Diabetol* 1: 4-9.
28. Jakubowicz D, Wainstein J, Ahren B, Landau Z, Bar-Dayana Y, et al. (2015) Fasting until noon triggers increased postprandial hyperglycemia and impaired insulin response after lunch and dinner in individuals with type 2 diabetes: a randomized clinical trial. *Diabetes Care* 38: 1820-1826.
29. Park YM, Heden TD, Liu Y, Nyhoff LM, Thyfault JP, et al. (2015) A high-protein breakfast induces greater insulin and glucose-dependent insulinotropic peptide responses to a subsequent lunch meal in individuals with type 2 diabetes. *J Nutr* 145: 452-458.
30. Yagi M, Kishimura Y, Okuda F, Ogura M, Ishizaki K, et al. (2018) Effect of yogurt on postprandial blood glucose after steamed rice intake. *Glycative stress res* 5: 68-74.
31. Jakubowicz D, Wainstein J, Landau Z, Ahren B, Barnea M, et al. (2017) High-energy breakfast based on whey protein reduces body weight, postprandial glycemia and HbA1C in Type 2 diabetes. *J Nutr Biochem* 49:1-7.
32. Kim Y, Keogh JB, Clifton PM (2016) Differential Effects of Red Meat/Refined Grain Diet and Dairy/Chicken/Nuts/Whole Grain Diet on Glucose, Insulin and Triglyceride in a Randomized Crossover Study. *Nutrients* 30: E687.
33. Dalla Man C, Micheletto F, Sathananthan M, Vella A, Cobelli C (2016) Model-Based Quantification of Glucagon-Like Peptide-1-Induced Potentiation of Insulin Secretion in Response to a Mixed Meal Challenge. *Diabetes Technol Ther* 18: 39-46.
34. Tricò D, Baldi S, Tulipani A, Frascerra S, Macedo MP, et al. (2015) Mechanisms through which a small protein and lipid preload improves glucose tolerance. *Diabetologia* 58: 2503-12.